

XX.- REFERENCIAS

- ❖ Albinson, T.F., 1989, Vetas mesotermales auríferas del sector norte del estado de Sonora, Memorias Técnicas, XVIII Convención Nacional de la Asociación de Ingenieros de Minas, Metalurgistas y Geólogos de Mexico, A.C., 20-40.

- ❖ Anderson, T. H., y Silver, T. H., 1977, U-Pb isotope ages of granitic plutons near Cananea: Sonora, Economic Geology and the Bulletin of the Society of Economic Geologists, 72(5): 827-836.

- ❖ Anderson, T.H., and Silver, L.T., 1979, The role the Mojave-Sonora megashear in the tectonic evolution of northern Sonora, in Anderson, T.H. and Roldán Quintana, J., eds, Geology of northern Sonora: Geological Society of America Annual Meeting, San Diego, California, Guidebook Field Trip 27, p. 59-68.

- ❖ Anderson, T.H., and Silver, L.T., 1981, An overview of Precambrian rocks in Sonora, México: Universidad Nacional Autónoma de México, Instituto de Geología, Revista, v. 5, no. 2, p. 131–139.

- ❖ Araux Sánchez E. 2000. Geología y Yacimientos Minerales de la sierra pinta, Municipio de Puerto Peñasco, Sonora; Universidad de Sonora. Tesis Maestría.

- ❖ Calmus, T., and M. Sosson, 1995, Southwestern extensión of the Pápago terrane into the Altar Desert region, northwestern Sonora, and its implications, in Studies on the Mesozoic of Sonora and Adjacent Areas, edited by C. Jacques-Ayala, C.M. Gonzalez-Leon, and J. Roldán-Quintana, Spec. Pap. Geol. Soc. Am. 301, 99-109.

- ❖ Campa, M. F., Coney, P. J., 1983. Tectono-stratigraphic terrenes and mineral resource distributions of México: Canadian Journal of Earth Sciences, 20, 1040-1051.

- ❖ Campbell, P.L., and Anderson, T.H., 2003, Structure and kinematics along a segment of the Mojave-Sonora megashear: A strike-slip fault that truncates the Jurassic continental magmatic arc of southwestern North America: Tectonics, v. 22, no. 6, p. 16-1–16-21.

- ❖ Coney, P.J., and S.J. Reynolds, 1977, Cordilleran Benioff zones, Nature, 270, 403-406.

- ❖ Cooper, C. A., y Arellano, A. R. V., 1946,Stratigraphy near Caborca, northwest Sonora, Mexico: Am. Ieum Geologists Bull., v. 30, p. 606-619.

- ❖ Damon, P.E.,Shafiqullah, M., and Clark, K.F. 1983 Geochronology of the porphyry copper deposits and related mineralization of Mexico; Canadian Journal of Earth Sciences; v.20, p.1052-1071.

- ❖ González-León, C.M., 1980, La Formación Antimonio (Triásico Superior-Jurásico Inferior) en la Sierra del Álamo, Estado de Sonora: Revista Instituto de Geología, Universidad Nacional Autónoma de México, 4, 13-18.

- ❖ González-León, C. and Jacques-Ayala. C. 1980. Paleogeografía del Cretácico temprano en Sonora. Actas de la facultad de Ciencias de la Tierra UANL. Linares 4. Pág. 125-152, edit Oct. 1990, Linares, México.

- ❖ Goldfarb R.J., Phillips G.N., Nokleberg W.J., 1998, Tectonic setting of synorogenic gold deposits of the Pacific Rim, Elsevier, Ore Geological Reviews 13, p 185-218.

- ❖ Goldfarb R.J., Groves D.I., Gardoll S, 2001, Orogenic gold and geologic time: a global synthesis, Elsevier, Ore Geological Reviews 18, p 1-75.

- ❖ González M.A. 1981. Cerros Gamuza y el Arpa. In. Longoria J., (editor). Regional geology of northwest Sonora. Geol. Soc. Amer. Cordilleran Section, Annual Meeting, Hermosillo, Sonora, México, Proceedings.

- ❖ González, L.C., Stanley, G.D. Jr., 1993. The Antimonio terrane of western Sonora and its paleogeographic significance, in Gonzalez, L.C. and Vega, G.L, eds. III simposio de la geología de Sonora y áreas adyacentes. Inst. de Geol. UNAM, UNISON. Pp. 41-43.

- ❖ Groves, D.I., Foster, R.P., 1991. Archean lode-gold deposits. In: Foster, R.P._Ed., Gold Metallogeny and Exploration. Blackie and Son, Glasgow, pp. 63–103.

- ❖ Groves D.I. et al., 1998, Orogenic gold deposits: A proposed classification in the context of their crustal distribution and relationship to other gold deposit types: Elsevier, Ore Geological Reviews 13, p 7-27.

- ❖ Harrar, N.G., 1989, Geology and paleomagnetism of the central Sierra El Batamote, northwestern Sonora, Mexico (M.S. Thesis); Cincinnati, OH, University of Cincinnati, 134p.

- ❖ Haxel, G.B., Tosdal, R.M., May, D.J., and Wright, J.E., 1984, Latest Cretaceous and early Tertiary orogenesis in south-central Arizona: Thrust faulting, regional metamorphism, and granitic plutonism: Geological Society of America Bulletin, v. 95, p. 631–653.

- ❖ Haxel, G.H., Wrigthy, J.E., May. D.J. y Tosdal, R.M., 1984. Reconnaissance geology of the Mesozoic and lower Cenozoic rocks of the southern Pápago Indian Reservation, Arizona: a preliminary report, in Jenney, J.P., and Stone, C. eds., Studies in western Arizona: Arizona Geol. Soc. Digest, v. 12, p. 17-29.

- ❖ Herrera-Urbina. Saúl y Pérez-Segura Efrén, 1990. Facies de metamorfismo en las secuencias de las inmediaciones del Poblado de Estación Llano, Sonora. Bol. Depto. Geol. Uni-Son, 1990. Vol. 1 y 2; Pág.; 7-28.

- ❖ Hodgson, C.J., 1993. Mesothermal lode-gold deposits. In: Kirkham, R.V., Sinclair, W.D., Thorpe, R.I., Duke, J.M. Eds., Mineral Deposit Modeling. Geol. Assoc. Can., Spec. Pap., 40, 635–678.

- ❖ INEGI, 1997. Carta Fisiográfica escala 1:1000,000.

- ❖ Iriondo, A., 2001, Proterozoic basements and their Laramide juxtaposition in northwestern Sonora, Mexico: Tectonic constraints on the southwestern margin of Laurentia [Ph.D. thesis]: Boulder, Colorado, University of Colorado, 222 p.

- ❖ Jacques-Ayala C. and Potter, P.E. 1987. Stratigraphy and Paleogeography of Lower Cretaceous rocks, Sierra El Chanate, Northwest, Sonora, México. (In) Dickinson W.R. and Klute. M.A. (Eds) Mesozoic rocks of Southern Arizona and adjacents areas. Arizona Geol. Soc. Digest. Pag; 203-214.

- ❖ Jaques-Ayala, Cesar; García y Barragán, J.C.; DeJong, K. A.; Graja;es-Nishimura, J. M.; López-Martínez, Margarita; and Layer, Paul, 1993, Age Constraints for Cretaceous-early Tertiary thrusting and faulting, northwestern Sonora: Universidad Nacional Autónoma de México, Instituto de Geología, y Universidad de Sonora, Departamento de Geología,

Simposio sobre la geología de Sonora y áreas adyacentes, 3rd,
Resúmenes. P. 61-63 (abstract).

- ❖ Kerrich, J. and Wyman, D. 1990. Geodynamic setting of mesothermal gold deposits an association with accretionary tectonic. *Geology*, v. 18, p. 882-885.
- ❖ Kerrich, R., 1993. Perspectives on genetic models for lode-gold deposits. *Miner. Deposita* 28, 362–365.
- ❖ López Martinez M. 1980, dato tomado de carta geológica Cananea 1:250000, COREMI.
- ❖ Longoria J.F., González M.A., Mendoza J.J., Pérez V.A. 1978. Consideraciones estructurales en el Cuadrángulo Pitiquito-La Primavera, NW Sonora., Bol. Depto Geol. Uni-Son, Hermosillo, Sonora, México, 1, pp. 119-144.
- ❖ LYNCH, D.J., 1981, Genesis and geochronology of alkaline volcanism in the Pinacate volcanic field, northwestern Sonora, Mexico: Ph.D. Thesis, Arizona University, 251 p.
- ❖ Martínez-Catalán José Ramón, 2002, Geología estructural y dinámica global: comportamiento dúctil; Universidad de Salamanca, Madrid España, pag. 197-268.

- ❖ Merriam, Richard, e Eells, J . L., 1979, Reconnaissance geologic map of the Caborca Quadrangle, Sonora, co: Univ. Sonora, Dept. Geología, Bol., v. 1, p. 87-94.
- ❖ Nesbitt, B.E., 1991. Phanerozoic gold deposits in tectonically active continental margins. In: Foster, R.P._Ed., Gold Metallogeny and Exploration. Blackie and Son, Glasgow, pp. 104–132.
- ❖ Noriega Martínez O. 2006 Evidencias de dos etapas de Mineralización de oro mesotermal en las área de Tajitos, Juárez-San Francisco, en la región de Caborca, Sonora México; Universidad de Sonora. Tesis de maestría.
- ❖ Nourse, J.A., 1995, Jurasic-Cretaceous paleogeography of the Magdalena region, northern Sonora, and its influence on the positioning of Tertiary metamorphic core complexes, in Jacques-Ayala, C., Gonzales León, C.M., Roldán Quintana, J., eds., Studies on the Mezosoic of Sonora and adjacent areas: Geological Society of America Special Paper 301, p. 59-78.
- ❖ Nourse, J.A., Iriondo, A., Premo, W.R., Kunk, M.J., and Stahl, E.R., 2000, New constraints on the southwestern margin of Laurentia from the Pinacate– Cabeza Prieta basement complex: Sonora and Arizona: Geological Society of America Abstracts with Programs, v. 32, no. 7, p. A-318.

- ❖ Pérez S. E. 1989. Los yacimientos de oro y plata de Sonora, México y sus relaciones con la Geología regional. En Delgado-Argote L.A. y Barajas, M. Eds. Contribuciones a la tectónica del oeste de México. Unión Geofísica Mexicana. Monografía No. 1. Pp. 147-174.

- ❖ Pérez-Segura, Cheilertz, Herrera-Urbina y Hanes (1996). Geología, mineralización, alteración hidrotermal y edad del yacimiento de oro de San Francisco, Sonora. Un depósito mesotermal en el noreste de México. Revista mexicana de Ciencias Geológicas, volumen 13, número 1996, p 65-89. Universidad Autónoma de México, Instituto de Geología, México, D.F.

- ❖ Robert, F., 1996. Quartz–carbonate vein gold. In: Eckstrand, O.R., Sinclair, W.D., Thorpe, R.I._Eds., Geology of Canadian Mineral Deposit Types. The Geology of North America, vol. P-1. Geol. Soc. Am., pp. 350–366.

- ❖ Shafiqullah, M., Damon, P., Lynch, D., Reynolds, S.J., Rehigh, W., Raymond, R., 1980, K-Ar geochronology and geologic history of southwestern Arizona and adjacent areas, *in* Jenney, J.P., Stone, C. (eds.), Studies in western Arizona: Tucson, Arizona Geological Society Digest, 12, 201-260.

- ❖ Silberman, L.M., Giles , D.A., Wenrich, K.J. and Morre-Hall A., 1987. Geochemistry of gneiss hosted gold deposits in northwestern Sonora, Mexico (abs.) Am. Inst: Mining Metall. Petroleum Engineers, Ann.

Meeting, 116th Denver, Colorado, Feb. 23-27, 1987. Abstracts with programs, p. 30.

- ❖ Silberman, M.L.; Giles, D.A.; y Graubard, 1988 Characteristics of gold deposits in northern Sonora, Mexico a preliminary report. *Economic Geology*, v.83, p.1966-1974.
- ❖ Silver, L.T., and Anderson, T.H., 1974, Possible left-lateral early to middle Mesozoic truncation of the southwestern North American craton: Geological Society of America Abstracts with Programs, v. 6, p. 955.
- ❖ Stewart, J. H, 1971. Basin and Range structure: a system of horsts and grabens produced by deep-seated extension: *Geological Society of America Bulletin*, v. 82, p. 1019-1044.
- ❖ Stewart, J. H. et al., 1986. Late Triassic paleogeography of the southern cordillera; the problem of a source for voluminous volcanic detritus in the Chinle Formation of the Colorado Plateau region, U.S. Geol.. Survey, p. 567-570, vol. 14 ; No. 7.
- ❖ Stewart J.H., McMenamin, M.A.S., Morales Ramírez, J.M. 1984. Upper Proterozoic and Cambrian rocks in the Caborca Region, Sonora, México. Physical stratigraphy, biostratigraphy, paleocurrent studies, and regional relations. U.S. Geol.. Survey Prof. Paper 1309. 36 p.

- ❖ Tosdal, R. M.; Haxel, G. B.; and Wright, J. E., 1989, Jurassic geology of the Sonoran Desert region, southern Arizona, southeastern California, and northernmost Sonora – construction of a continental-margin magmatic arc, in Jenney, J. P., and Reynolds, S. J., eds., Geologic evolution of Arizona: Arizona Geologiclas Society Digest, v. 17, p. 397-434.

- ❖ Thompson, T.B. (ed.), 1998, Gold deposits of Northern Sonora, Mexico: Society of Economic Geologists, Guidebook Series, Vol. 30, 252 p.