

VIII. BIBLIOGRAFIA

Al-Aswad M.B. (1971). The amino acids of some Iraqi dates. *Journal Food Science*, 36: 1019-1020.

Allaith, A.A.A. (2008). Antioxidant activity of Bahraini date palm (*Phoenix dactylifera* L.) fruit of various cultivars. *International Journal Food Science and Technology*, 43, 1033-1040

Al-Farsi, M., Alasalvar C.; Morris A, Baron M.; Shahidi F. (2005a). Composition and Sensory Characteristics of Three Native Sun-Dried Date (*Phoenix dactylifera* L.) Varieties Grown in Oman. *Journal Agriculture Food Chemistry* 53, 7586-7591.

Al-Farsi, M.; Alasalvar, C.; Morris, A.; Baron, M.; Shahidi, F. (2005b). Comparison of Antioxidant Activity, Anthocyanins, Carotenoids, and Phenolics of Three Native Fresh and Sun-Dried Date (*Phoenix dactylifera* L.) Varieties Grown in Oman. *Journal Agriculture Food Chemistry* 53, 7592-7599.

Al-Farsi, M. y Lee, C. (2008). Nutritional and Functional Properties of Dates: A Review. *Critical Reviews in Food Science and Nutrition*, 48: 877-887.

Al-Hooti S.; Sidhu J.; Al-Otaibi J.; Al-Ameeri H.; Qabazard H. (1997). Date bars fortified with almonds, sesame seeds, oat flakes and skim milk powder. *Plant Foods for Human Nutrition* 51:125-135.

Al-Hooti S.; Sidhu J.; Qabazard H. (1997). Physicochemical characteristics of five date fruit cultivars grown in the United Arab Emirates. *Plant Foods for Human Nutrition* 50:101-113.

Al-Khayri J.M. (2005). Date Palm *Phoenix dactylifera* L. Protocol for Somatic Embryogenesis in Woody Plants, editado S.M. Jain y P.K. Gupta 309-319.

Al-Shahib, W. y Marshall, R.J. (2002). Dietary fibre content of dates from 13 varieties of date palm *Phoenix dactylifera* L. *International Journal Food Science and Technology* 37, 719-721.

Aguilar, M.M.A. (2005). Propiedades físicas y mecánicas de las películas biodegradables y su empleo en el recubrimiento de frutos de aguacate. Tesis de Maestría, Instituto Politécnico Nacional.

A.O.A.C. (2000). The Official Methods of Analysis of the Association of Official Analysis Chemistry. Vol II. 17th Edition. Dr. W. Horwitz Editor, Gaithersburg, Maryland

Arizmendi, C.D. y Domínguez, L.A. (2009). Optimización de dos compuestos plastificantes (glicerol y polietilenglicol) en la elaboración de una película plástica comestible obtenida a partir del mucílago de nopal de la especie "*Opuntia tomentosa* Salm-Dyck". Universidad Autónoma del Estado de México

Auda, H.; Al-Wandawi, H.; Al-Adhami, L. (1976). Protein and Amino Acid Composition of Three Varieties of Iraqi Dates at Different Stage of Development. *Journal Agriculture Food Chemistry*, 24, 365-367.

Ayala-Chavez, G.M. (2009). Establecimiento de las Condiciones de Proceso para dar Valor Agregado a la Producción de Nuez (*Carya illinoensis* L.) e Higo (*Ficus carica* L.) en el Estado de Sonora. Tesis de Maestría, Centro de Investigación en Alimentación y Desarrollo, A.C. Hermosillo, Sonora, México.

Azeredo, H.; Mattoso L.; Wood D.; Williams T.; Avena-Bustillos R.; McHugh T. (2009). Nanocomposite Edible Films from Mango puree Reinforced with Cellulose Nanofibers. *Journal of Food Science*, Vol. 74, Nr. 5.

Badui, D.S. (1999). Química de los Alimentos. 4° ed. México. Longman de México Editores S.A. de C.V.

Baldwin, E.; Nisperos-Carriedo, M.; Baker, R. (1995). Edible Coating for Lightly Processed Fruits and Vegetables. *HortScience*, Vol. 30(1), pág. 35-38, Febrero.

Baldwin, E.; Nisperos-Carriedo, M.; Hagenmaier, R.; Baker, R. (1997). Using Lipids in Coatings for Food Products. *Food Technology*. Vol. 64, pág.. 56-61.

Barreveld, W.H. (1993). Date palm products. *Agricultural Services Bulletin* 101; FAO, Rome, Italy.

Beer, F.; Johnston, E.; DeWolf, J.; Mazurek, D. (2010). Mecánica de materiales 5ta. edición, editorial McGraw-Hill, pág. 54, México.

Brindle, L.P. y Krochta, J.M. (2008). Physical Properties of Whey Protein-Hydroxypropylmethylcellulose Blend Edible Films. *Journal of Food Science*. Vol. 23. Nr 9, pág. 446-454.

Buitimea A.C. (2005). Extracción y caracterización fisicoquímicas de pectinas del nopal. Tesis de Licenciatura. Universidad Juárez Autónoma de Tabasco, Extensión de los Ríos. Tabasco, México, pág. 91.

Blumenkrantz N. y Asboe-Hansen G. (1973). New Method for Quantitative Determination of Uronic Acids. *Analytical Biochemistry* 54, pág. 484-489.

Byars, E., Snyder R. (1978). Mecánica de cuerpos deformables, 3ra edición, Representaciones y servicios de ingeniería, S.A. México, pág. 438.

Constenla D. y Lozano J.E. (2003). Kinetic model of pectina demethylation. Latin American Applied Research, Volumen 33, número 2.

Chaira, N; Ferchichi, A.; Mrabet, A.; Sghairoun, M. (2007). Chemical Composition of the Flesh and the Pit of Date Palm Fruit and Radical Scavenging Activity of Their Extracts. Pakistan Journal of Biological Sciences 10(13):2202-2207.

Chambi, H. y Grosso, C. (2006). Edible films produced with gelatin and casein cross-linked with transglutaminase. Food Research International, 39, pág. 458-466.

Copley, M.; Rose P.; Clapham, A.; Edwards, D.; Horton M.; Evershed, R. (2001). Proceedings of the Royal Society of London. 268, 593-597.

Debeaufort, F; Quezada-Gallo, J.; Voilley, A. (1998). Edible Films and Coatings: Tomorrow's Packagings: A Review. Critical Reviews in Food Science, 38 (4): 299-313.

Devia-Pineda, J.E. (2003). Proceso para producir pecinas críticas. Revista Universidad EAFIT, Número 129, pág. 21-30.

Diab, T.; Biliaderis, C.; Gerasopoulos, D.; Svangelos, E. (2001). Physicochemical properties and application of pullulan edible films and coatings in fruit preservation. Journal of the Science of Food and Agriculture 81:988-1000.

Díaz-Sánchez, D.D. (2008). Elaboración de una barra multigrano con relleno de mermelada de higo (*Ficus Carica L.*) variedad Kadota. Tesis de Maestría. Centro de Investigación en Alimentación y Desarrollo A.C., Hermosillo, Sonora

El-Mardi, M.; Esechie, H.; Al-Kharousi, L.; Abdelbasit, K.M. (2002). Effect of Pollination Method on Changes in Physical and Chemical Characteristics of Date Fruit During Development. Agricultural Sciences, 7 (1):21-27.

El-Zoghbi M. (1994). Biochemical changes in some tropical fruits during ripening. Food Chemistry 49, 33-37.

Eltayeb, E.; Al-Hasni A.; Farooq, S. (1999). Changes in Soluble Sugar Content During the Development of Fruits in Some Varieties of Oman Date Palm (*Phoenix dactylifera*) Pakistan Journal of Biological Sciences 2(1):255-258.

Falade K.O. y Abbo E.S. (2007). Air-drying and rehydration characteristics of date palm (*Phoenix dactylifera* L.) fruits. *Journal of Food Engineering* 79:724-730.

Food Agriculture Organization (FAO). (2007). Promoción de la producción y mejoramiento de dátiles. TCP/MEX/3102(D).

Faostat. (2011). Producción de productos alimenticios y agrícolas. País por producto, dátiles en 2009. <http://faostat.fao.org/DesktopDefault.aspx?PageID=339&lang=es> Revisado en 2011.

Gennadios A. y Weller C. (1990). Edible Films and Coating from Wheat and Corn Proteins. *Food technology*, pág. 63-69.

Gennadios, A., Weller, C.L., y Gooding, C.H. (1994). Measurement Errors in Water Vapor Permeability of Highly Permeable, Hydrophilic Edible Films. *Journal Food Eng.* pág.395-409.

Golshan, T. y Fooladi, M. (2005). Changes in Physical and Chemical Characteristic of Mozafati Date Fruit During Development. *Journal of Biological Sciences* 5 (3): 319-322.

Goycoolea, V. y Cárdenas, B. (2003). Pectin from *Opuntia* spp.: A Short Review. *Journal of the Professionals Association of Cactus Development*.5:17-29.

Hernandez-izquierdo, V. y Krochta, J.M. (2008). Thermoplastic Processing of Proteins for Films Formation- A Review. *Journal of Food Science*, Vol. 73, Nr,2.

Hui Y.H. (2006). *Handbook of Fruits and Fruit Processing*, Blackwell Publishing, pág. 189-193.

Ihsanullah,I.; Iqbal,Y.; Khattak, N. (2005).Effect of various irradiation doses on some nutrients of Pakistani date. *Journal of Radioanalytical and Nuclear Chemistry*, Vol. 266, No. 2, pág. 361-366.

Ishurd, O.; Zahid, M.; Xiao, P.; Pan Y. (2004). Protein and amino acids contents of Libyan dates at three stages of development. *Journal of the Science of Food and Agriculture*. Volumen 84, Issue 5, pág. 481–484.

International Pectin Producers Association (IPPA).(2011). Dirección en internet http://www.ippa.info/what_is_pectin.htm

Jaradat A y Zaid A. (2004). Quality traits of date palm fruits in a center of origin and center of diversity. *Food, Agriculture & Environment* Vol.2 (1):208-217.

Kemper, S.L. y Fennema, O.R. (1984) a. Water Vapor Permeability of Edible Bilayer Films. *J. Food Sci.* pág. 1478-1481.

Kemper, S.L. y Fennema, O. R. (1984) b. Water Vapor Permeability of an Edible, Fatty Acid Bilayer Film. *J. Food Sci.* pág. 1482-1485.

Kester, J.J. y Fennema, O.R. (1986). Edible films and coatings: a review. *Food technology*, pág. 47-59.

Krochta, J. y De Mulder-Johnston, C. (1997). Edible and biodegradable polymer films: Challenges and opportunities. *Food technology*, pág. 61-74.

Kwaasi, A.A.; Harfi, H.A.; Parhar, R.S.; Al-Sedairy, S.T.; Collison, K.S.; Panzani, R.C; Al-Mohanna, F.A. (1999). Allergy to date fruits: characterization of antigens and allergens of fruits of the date palm (*Phoenix dactylifera* L.). *Allergy*, 54, pág. 1270-1277.

Lin S.Y. y Krochta J.M. (2006). Whey protein coating efficiency on mechanically roughened hydrophobic peanut surfaces. *Journal of Food Science*, 71 (6):270-275.

Lin, D. y Zhao Y. (2007). Innovation in the Development and Application of Edible Coatings for Fresh and Minimally Processed Fruits and Vegetables. *Institute of Food Technology*, pág. 60-75.

Llanos-Navarro, María. (2007). Efecto de la composición de recubrimientos comestibles a base de hidroxipropilmetilcelulosa y cera de abeja en la calidad de ciruelas, naranjas y mandarinas. Tesis de Doctorado. Universidad Politécnica de Valencia. Valencia, España.

Martin-Polo, M; Mauguin, C.; Voilley, A. (1992). Hydrophobic Films and Their Efficiency Against Moisture Transfer. 1. Influence of the Film Preparation Technique. *Journal of Agriculture and Food Chemistry*, 40 (3), pág. 407–412.

Matsuhiro, B. y Rubio, M.J. (2001). Modificación química de pectina de limón. *Boletín de la Sociedad Chilena de Química*. Volumen 46, número 4.

Mazlounzadeh, S.; Shamsi, M.; Nezamabadi-Pou, H. (2010). Fuzzy logic to classify date palm trees based on some physical properties related to precision agriculture. *Precision Agriculture* 11:258-273.

McHugh, T.H. y Krochta, J. (1994). Water vapor permeability properties of edible whey protein-lipid emulsion films. *Journal American Oil Chemistry Society* 71:307-312.

McHugh, T.H.; Huxsoll C.; Krochta J. (1996). Permeability properties of fruit puree edible films. *Journal Food Science*. 61:88-91.

Mc Hugh, T.H. y Senesi, E. (1996). Apple Wraps: A Novel Method to Improve the Quality and Extend the Shelf Life of Fresh-cut Apples. *Journal of Food Science*, Vol. 65, No. 3.

Meza Godoy, A. (2006). Desarrollo de películas o recubrimientos comestibles con potencial para el recubrimiento de frutas frescas. Universidad Autónoma Metropolitana-Iztapalapa.

Murrieta-Becerra, María Luisa. (2010). Caracterización de Higo (*Ficus carica* L.), Variedad Kadota y su Potencial como Película Comestible. Tesis de licenciatura Instituto Tecnológico del Valle del Yaqui, Hermosillo, Sonora, México.

Mustafa, A.; Harper, D.; Johnson, D. (1986). Biochemical Changes During Ripening of Some Sudanese Date Varieties. *Journal of Science Food and Agriculture*, 37,pág. 43-53.

Myhara, R., A. Al-Alawi, J. Karkalas, M.S Taylor. (2000). Sensory and textual changes in maturing Omani dates. *Journal Science Food and Agriculture* 80:2181-2185.

Myhara, R; Karkalas, J.; Taylor, M. (1999). The composition of maturing Omani dates. *Journal of the Science Food and Agriculture*,79:1345-1350.

Nisperos-Carriedo, M.O. (1994). Edible Coatings and Films Based on Polysaccharides. En: *Edible coatings and Films to Improve Food Quality*. J.M. Krochta, E.A. Baldwin y M. Nisperos-Carriedo (Eds.) Technomic, Lancaster, Pensilvania, Estados Unidos de América, pág. 305-336.

Prosky, L., Asp., N.-G., Schweizer, T.F., DeVries, J.W. and Furda,I. (1988). Determination of insoluble, soluble, and total dietary fibre in foods and food products. Interlaboratory study. *J. Assoc. Off. Anal. Chem.* 71: 1017-1023.

Rojas-Graü, M. (2005). Informe de la estancia realizada en el Departamento de Agricultura de los Estados Unidos (USDA). California, Estados Unidos de América.

Rojas-Graü, M.; Avena-Bustillos, R.; Friedman, M.; Henika, P.; Martín-Belloso, O.; McHugh TH. (2006) .Mechanical barrier and antimicrobial properties of Apple puree edible films containing plant essential oils. *Journal of Agriculture and Food Chemistry* 54 (24):9262-9267.

Roy, S.K.; Khurdiya, D.S.; Waskar, D.P.; (1995). Capítulo 28, Other Subtropical Fruits. *Handbook of Fruit Science and Technology: production, composition, storage and processing*. Edited by D.K. Salunkhe y S.S. Kadam, 539-346.

SAGARPA. Secretaría de Agricultura Ganadería, Desarrollo Rural, Pesca y Alimentación .Servicio de información agroalimentaria y pesquera. Cierre de la producción agrícola por estado del año 2010. Liga http://www.siap.gob.mx/index.php?option=com_wrapper&view=wrapper&Itemid=351 (2011).

Sánchez D.; Aguilar, C.N.; Contreras J.C.; Nevarez, G.V. Moléculas pécticas: extracción y su potencial aplicación como empaque. *Tecnociencia Chihuahua*. Volumen 5, No. 2. 2011

Sawaya, W.N.; Khatchadourian, H.A.; Khalil, J.K. ; Safi, W.M. ; Al-Shalhat, A. (1982). Growth and compositional changes during various developmental stage of Saudi Arabian date cultivars. *Journal Food Science* 47:1489.

Sawaya, W.N.; J. Khalil, and W.J. Safi, (1984). Chemical Composition and Nutritional Quality of Date Seeds. *Journal Food Science* 49:617.

Shidhu, J.S.; Al-Saqer, J.M.; Al-Hooti, S.; Al-Othman, A. (2003). Quality of Pan Bread Made by Replacing Sucrose With Date Syrup Produced by Using Pectinase/Cellulase Enzymes. *Plant Food for Human Nutrition* 58: 1-8.

Sidhu, J.S. (2006). Capítulo 22 .Date fruits Production and Processing. *Handbook of fruits and fruits processing*. Edited by Y.H. Hui Pp 391-419.

The National Academies Press. (2005). Capítulo 7 .Dietary, Functional and total fiber. *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids*. Washington, D.C.:Autor.

USDA 2011. United States Department of Agriculture. Liga en internet: <http://ndb.nal.usda.gov/ndb/foods/show/2519>

Vayalil, P.K. 2002. Antioxidant and Antimutagenic properties of aqueous extract of date palm (*Phoenix dactylifera* L. *Arecaceae*) *Journal Agriculture Food Chemistry* 50,610-617.

Villada ,H.S.; Acosta, H.A.; Velasco, R.J. (2007). Biopolímeros naturales en empaques biodegradables. *Temas Agrarios*, 12(2):5-13.

Villagómez-Zavala, D.; Gómez-Corona, C.; San Martín, E.; Pérez-Orozco, J.; Vernon-Carter, E.; Pedroza-Islas, R. (2008). Comparative study of the mechanical properties of edible films made from single and blended hydrophilic biopolymer matrices. *Revista Mexicana de Ingeniería Química*. Vol. 7, No. 3, pág. 263-273.

Vithanage, C.R.; Grimson, M.J.; Wills, P.; Harrison, P.; Smith, B.G. (2010). Rheological and structural properties of high-methoxyl esterified, low-methoxyl esterified and low-methoxyl amidated pectin gels. *Journal of Texture Studies*, 41: 899-927.

Vinson J.A. (1999). The functional food properties of figs. *Cereal Foods World*, pág. 44, 82-87.

World Health Organization & Food Agriculture Organization (WHO/FAO). (2003). A report of joint WHO/FAO expert consultation on diet, nutrition and the prevention of chronic diseases, Technical Report Series No.916, Geneva, Switzerland.

www.a-campo.com.ar, (2011). Liga directa en internet: www.a-campo.com.ar/espanol/fruticultura/fruticult10.htm

Zemansky, A.C. y Dittman, M.M. (1981). *Calor y termodinámica*. 6ta ed. Libros Mc Graw-Hall de México, pág. 228.

