

BIBLIOGRAFIA

Abbey, B.W. (1985). Food uses of whole oil and protein seed (1989). By the American Oil Chemists Society. Edited by Edmund W. Lusas, David R. Erickson and Wai Kit Nip.

(AOAC) Association of Official Analytical Chemist (1990). Official Methods of Analysis. 15 Th. Edition.

Altschyl M. A. (1974). New Protein Foods. Ed. Academy Press EUA. Vol. 1 Technology. pág. 249.

(ASTM) Standards for Bomb Calorimetry and Combustion Methods (1977). American Society for Testing and Materials.

Aykroyd, W.R. and J.J. Doughthy (1964). Legumes in Human Nutrition. F.A.O. Food J. Nutrition. Rome.

Baldi, G. y F. Salamini (1973). Variability of Essential Amino Acid content in seed of 22 Phaseolus species, Theoretical and Applied Genetics. Vol. 43: 75 - 78.

Bidwell, R.G.S. (1987). Fisiología vegetal. 1a. edición, AGT editor, pág. 207 y 208.

Bouscaren, S.J., J.G. Waines y L.A. Boykm-Bouscaren (1983). El cultivo y uso de los Téparis en Sonora, México. Potencial del frijol Tépari en las zonas desérticas, Traducción de la revista Desert Plants. Vol. 5 (1).

Brand J.C. and V. Cherikoff (1985). The Nutritional Composition of Australian Aboriginal Food Plants of the Desert Regions, In: Plant for Arid Lands, Wickens G. E., J.R. Goodin and D.V. Field. 1a. Ed., pág. 55 y 64.

Bressani, R. (1985). Nutritive Value of Cowpea. In: Cowpea Research, Production and Utilization. Singh S.R. and K.O. Rachie. Ed. John Wiley & Sons, Gran Bretaña, pág. 356 y 357.

Bressani, R. and L.G. Elias (1978). Nutritional Value of Legume Crops for Humans and Animals. (INCAP). Guatemala. (A).

Bressani, R. and L.G. Elias. (1978). Seed Protein Improvement in Cereals and Grain Legumes. The World Protein and Nutritional Situation. pág. 18,19. (B).

Bressani, R. and L.G. Elias (1980). Nutritional Value of legume Crops for Humans and Animals. In:Advances in Legume Science. Summerfield, R.J. and A. H. Bunting. Eds.,Royal Botanic Gardens, Kew, pág. 137-147.

Bressani, R. and N.S. Scrimshaw (1961). The Development of INCAP Vegetable Mixtures, 1: Basic Animal Studies. In National Academy of Sciences (NAS), Meeting Protein Needs of Infants and Preschool Children. Washington, DC, NAS, Publication 843.

Buhrow, R. (1983). The Wild Bean of Southwestern North America, Desert Plants. Vol. 5(2): 67-88.

Cabezas, M.T., B. Cuévas., B. Murillo, L.G. Elías and R. Bressani(1982a). Evaluación Nutricional de la Sustitución de la Harina de Soya y Sorgo por Harina de Frijol Caupí Crudo (*Vigna sinensis*). Archivos Latinoamericanos de

Nutrición. 32, 559-578.

Cabezas, M.T., J. García, B. Murillo, L.G. Elías and R. Bressani (1982b). Valor Nutritivo del Frijol Caupí Crudo y Procesado. Archivos Latinoamericanos de Nutrición. 32, 543-558.

Carpenter, K. (1981). The Nutritional Contribution of Dry Beans (*Phaseolus vulgaris*) in Perspective. Food Technology. Vol. 35:77.

Cathey, H.M. y Campbell, L.E. (1987). Respuesta de la Planta a la Calidad y Cantidad de Luz. En: Mejoramiento de Plantas en Ambientes Poco Favorables. Christiansen, N.M. y Ch. F. Lewis. 1a. Ed., Editorial Limusa, México, pág. 23.

Chang, R., S. Schwimmer, and H.K. Burr (1977). Phytate Removal from Whole Dry Beans by Enzymatic Hydrolisis and Diffusion. J. Food Sci. Vol. 42: 1098-1101.

Del Valle, F.R., M. Escobedo, M.J. Muñoz, R. Ortega and H.J. Bourges. (1983). Chemical and Nutritional Studies on Mesquite Beans (*Prosopis juliflora*). J. of Food Sci. Vol. 48: 914.

Downie, N.M. y R.W. Heath (1975). Métodos Estadísticos Aplicados. Editorial HARLA, México, pág. 336.

Duke, J.A. (1981). Handbook of Legumes of World Economic Importance. Plenum Press. USA.

Earle, F.R. and Q. Jones (1960). Analyses of Seed Samples from 113 Plant Families. Economic Botany. June.

- Elston, J. and A.H. Bunting (1980). Water Relations of Legume Crops. In: Advances in Legume Science. Summerfield, R.J. and A.H. Bunting. Eds. Royal Botanic Gardens, Kew, pág. 37-41.
- El Tinay, A.H., S.O. Mahgoub, B.E. Mohamed and M.A. Hamed (1989). Proximate Composition and minerals and phytate contents of Legumes Grown in Sudan. Journal of Food Composition and Analysis. Vol. 2: 69-78.
- Ershow, A.G. and K. Wong-Chen (1990). Chinese Food Composition Tables. Journal of Composition and Analysis. Academic Press, Inc. USA. Vol. 3, Numbers 3/4 Sep/Dec.
- Esh, G.C., T.W. De and K.P. Basu (1959). Influence of Genetic Strain and environment on Protein Content of Pulses. Science. Vol. 129: 148-149.
- Federici, C.T., B. Ehdaie and J.G. Waines (1990). Domesticated and Wild Tepary Bean: Field Performance with and without Drought-Stress. Agronomy Journal. Vol. 82: 896-899.
- F.A.O. (1970). Contents of protein in Legumes Food.
- Felger, S.R. and P.G. Nabhan (1978). Agroecosystem Diversity: A model from the Sonoran Desert. In: Social and Technological Management in Dry Lands. Past and Present. Indigenous and Imposed. Gonzalez N. L. (ed.). AAAS, Selected Symposium #10, pág. 129-148.
- Figueiredo, A.A. (1990). Mesquite: History Composition, and Food Uses. Food Technology. Vol.44 (11) :118-128.
- Fitter, A.H. and R.K.M. Hay (1987). Environmental Physiology of plants. 2a. Edición, Academic, Inglaterra, pág. 130.

Freeman, G.E. (1912). Southwestern Beans and Teparies. Arizona Agr.. Exp. Sta. Bulletin 68.

Gepts ,P. and D. Debouck (1991). Origin, Domestication and Evolution of the Common Bean (*Phaseolus vulgaris L.*) Common beans: Research for crop improvement. CIAT Redwood Press Ltd., England, pág. 9.

Hughes, S. J. (1991). Potential Contribution of Dry Bean Dietary Fiber to Health. Food Technology. Vol.45 (9): 122.

Huterwal G.O. (1977). Hidroponia Cultivo de plantas sin Tierra, Ed. Albatros, Argentina. pág. 47.

Kadam S.S., S.S. Deshpande and N.D. Jambhale (1989). Seed Structure and Composition.In: C.R.C. Handbook of World Food Legumes: Nutritional Chemistry, Processing Technology and Utilization. Salunkhe D.K. and S.S. Kadam. Press EVA. Vol. 1: 38-45.

Koehler, H.H. and D.W. Burke (1988). Nutrient and Sensory Properties of Dry Beans (*Phaseolus vulgaris L.*) Grown Under Various Cultural Conditions. J. of Food Science. 53(4): 1135-1138.

Kohashi, S.J. (1990). Aspectos de la morfología y fisiología del frijol Phaseolus vulgaris L. y su relación con el rendimiento. 1a. Edición. Centro de Botánica. Colegio de Postgraduados, Chapingo-Montecillo, México, pág. 11,14 ,15, 38 y 39.

Larcher, W. (1980). Physiological Plant Ecology, 2a. Edición, Springer-Verlag, Alemania, pág. 1-3, 18, 206.

Liener, I.E. (1983). Toxic Constituents In Legumes.In: Chemestry and Biochemestry of legumes. Arora, S. K. (Editor). Edward Arnold. London.

Maron, S. H. y C. F. Prutton (1984). Fundamentos de Fisicoquímica. Decimacuarta Reimpresión, Editorial Limusa. pág. 143,144 y 151.

Mc Daniel, G.R. (1987). La fisiología de los efectos de la temperatura en las plantas.En: Mejoramiento de plantas en ambientes poco favorables. Christiansen, N.M. y Ch. F. Lewis. 1a. Edición, Editorial Limusa, México, pág. 23.

Meiners, C.R. , N.L. Derise, H.C. Lau, M.G. Crews, S.J. Ritchey and E.W. Murphy (1976). The Content of Nine Mineral Elements in Raw, Cooked Mature Dry Legumes. J. Food Chem. Vol. 24(6):1126-1130.

Miranda C. S. (1979). Evolución de *Phaseolus vulgaris* y *Phaseolus coccineus* in Engleman, E.M. (ed.). Contribuciones al conocimiento del frijol (Phaseolus) en México. Colegio de Postgraduados Chapingo, México, pág. 83-99.

Nabhan, G.P.,J.Berry and Ch. Weber (1979). Legumes in the Papago-Pima India Diet and Ecological Niche. Kiva. 44: 173.

Nabhan, G.P., J.W. Berry, C. Anson and C.W. Weber (1980). Papago Indian Floodwater Fields and Tepary Bean Protein Yields. Col. Food Nutr. 10:71

Nabhan, G.P. (1983). The Desert, Tepary as a Food Resource. Desert Plants. pág. 11-49.

Nabhan, G.P., C.W. Weber and J.W. Berry (1985). Variation in composition of Hopi Indian Beans. Ecology of Food and Nutrition. Vol.16:135-152.

National Academy of Sciences (1979). Tropical Legumes: Resources for the Future. National Academy of Sciences, Washington, D.C., pág. 305-307.

Norton, G., F.A. Bliss and R. Bressani (1985). Biochemical and Nutritional Attributes of Grain Legumes.In: Grain Legume Crops. Edited by R.J. Summerfield and E.H. Robertis.

O'Dell,B.L. and A.D. Boland (1976). Complexation of Phytate with Proteins and Cations in Corn Germand Oilseed Meals. J Agric. Food Chem. Vol. 24: 4-7.

Onayemi, O., W.G. Pond and L. Krook (1976). Effect of Processing on the Nutritive value of Cowpeas for the growing rat. Nutr. Reports. Int. 13 (3):29.

Pate, J.S. and F.R. Minchin (1980). Comparative Studies of Carbon and Nitrogen Nutrition of Selected Grain Legumes.In: Advances in Legume Science. Summerfield, R.J. and A.H. Bunting. Eds. Royal Botanic Gardens, Kew, pág. 105-113.

Peterkin, B., J. Nichols and C. Cromwell (1975). U.S. Department of Agriculture. Agriculture Information Bulletin No. 382.

Saeed, A.R. (1977). Grain Legumes in Sudan, Expert Consultation on Grain Legumes by Processing. Central Food Technological Institute, Mysore, India, Vol. 14: 18-24.

- Shreve, F. and I.L. Wiggins (1986). Vegetation and Flora of the Sonoran Desert. Stanford University Press, EUA, Vol. I: 143-145, 628-631, 729.
- Siegel, S. (1976). Estadística no Paramétrica: Aplicada a las Ciencias de la Conducta. Tercera reimpresión. Ed. Trillas. México. pág. 7, 143-148 y 305-309.
- Signoret, P. J. (1970). Datos sobre algunas características del mezquite (*Prosopis leavigata*), y su aprovechamiento en el Valle del Mezquital. En Mezquites Huisaches. IMRNR México. pág. 72-146.
- Sinclair, T.R. and C.T. De Wit (1975). Photosynthate and Nitrogen Requirements for Seed Production by Various Crops. Science. Vol.189: 566.
- Sotelo, A. (1981). Leguminosas silvestres, reserva de proteínas para la alimentación del futuro. Información Científica y Tecnológica. Vol. 3(54): 28.
- Steele, V.M. and K.L. Mehra (1980). Structure, Evolution, and Adaptation to Farming Systems and Environments in Vigna. In:Advances in Legume Science. Summerfield R.J. and A.H. Bunting . Eds. Royal Botanic Gardens, Kew, Inglaterra, pág. 393-403.
- Summerfield, R.J. and H.C. Wien (1980). Effects of Photoperiod and Air Temperature on Growth and Yield of Economic Legumes. In:Advances in Legume Science. Summerfield, R.J. and A.H. Bunting. Eds. Royal Botanic Gardens, Kew, pág. 17-33.
- Thorn, K.A., A.M. Tinsley, C.W. Weber and J.W. Berry . (1983). Antinutritional Factors in Legumes of the

Sonoran Desert. Ecol. Food and Nutrition. Vol.13: 251-256.

U. S. Department of Agriculture (1975). Composition of Foods.
Raw, Processed, Prepared. Agriculture Handbook No. 8.
Agricultural Research Service. pág. 9 y 10.

Ville, A.C. (1981). Biología. 7a. Edición. Editorial Interamericana.
México. pág. 112.

Waines G.J. (1978). Protein Contents, Grain Weights, and
Breeding Potential of Wild and Domesticated Tepary Beans.
Crop Science. Vol. 18: 587-589.

Wayne W. D. (1980). Bioestadística: Base Para el Análisis de las
Ciencias de la Salud. 2a. Reimpresión. Ed. Limusa. México.
pág. 377-387 y 454.